

The calculator for the olivine-orthopyroxene-spinel oxybarometer is built upon the ThermoEngine platform developed by ENKI (and Mark Ghiorso).

Coogan L. A., Saunders A. D., and Wilson R. N. 2014. Aluminum-in-olivine thermometry of primitive basalts: Evidence of an anomalously hot mantle source for large igneous provinces. *Chemical Geology* 368:1–10.

Jianping, L., Kornprobst, J., Vielzeuf, D., Fabriès, J., 1995. An improved experimental calibration of the olivine-spinel geothermometer. *Chin. J. of Geochem.* 14, 68–77.
<https://doi.org/10.1007/BF02840385>

Sack, R.O., Ghiorso, M.S., 1994a. Thermodynamics of multicomponent pyroxenes: I. Formulation of a general model. *Contr. Mineral. and Petrol.* 116, 277–286.
<https://doi.org/10.1007/BF00306497>

Sack, R.O., Ghiorso, M.S., 1994b. Thermodynamics of multicomponent pyroxenes: II. Phase relations in the quadrilateral. *Contr. Mineral. and Petrol.* 116, 287–300.
<https://doi.org/10.1007/BF00306498>

Sack, R.O., Ghiorso, M.S., 1994c. Thermodynamics of multicomponent pyroxenes: III. Calibration of $\text{Fe}^{2+}(\text{Mg})$ -1, $\text{TiAl}_2(\text{MgSi}_2)$ -1, $\text{TiFe}^{23+}(\text{MgSi}_2)$ -1, $\text{AlFe}^{3+}(\text{MgSi})$ -1, $\text{NaAl}(\text{CaMg})$ -1, $\text{Al}_2(\text{MgSi})$ -1 and $\text{Ca}(\text{Mg})$ -1 exchange reactions between pyroxenes and silicate melts. *Contr. Mineral. and Petrol.* 118, 271–296. <https://doi.org/10.1007/BF00306648>

Sack, R.O., Ghiorso, M.S., 1991a. Chromian spinels as petrogenetic indicators: Thermodynamics and petrological applications. *American Mineralogist* 76, 827–847.

Sack, R.O., Ghiorso, M.S., 1991b. An internally consistent model for the thermodynamic properties of Fe–Mg-titanomagnetite-aluminate spinels. *Contr. Mineral. and Petrol.* 106, 474–505. <https://doi.org/10.1007/BF00321989>

Wan Z., Coogan L. A., and Canil D. 2008. Experimental calibration of aluminum partitioning between olivine and spinel as a geothermometer. *American Mineralogist* 93:1142–1147.